

Common subspaces of L_p -spaces

ALEXANDER KOLDOBSKY

DEPARTMENT OF MATHEMATICS

UNIVERSITY OF MISSOURI-COLUMBIA

COLUMBIA, MO 65211

Abstract. For $n \geq 2, p < 2$ and $q > 2$, does there exist an n -dimensional Banach space different from Hilbert spaces which is isometric to subspaces of both L_p and L_q ? Generalizing the construction from the paper "Zonoids whose polars are zonoids" by R. Schneider we give examples of such spaces. Moreover, for any compact subset Q of $(0, \infty) \setminus \{2k, k \in \mathbb{N}\}$, we can construct a space isometric to subspaces of L_q for all $q \in Q$ simultaneously.

AMS classification: Primary 46B04, Secondary 46E30

Key words: isometries, positive definite functions, spherical harmonics.

e-mail: mathak@mizzou1.missouri.edu

Common subspaces of L_p -spaces

ALEXANDER KOLDOBSKY

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF MISSOURI-COLUMBIA
COLUMBIA, MO 65211

1. Introduction.

This work started with the following question: For given $n \geq 2, p \in (0, 2)$ and $q > 2$, does there exist an n -dimensional Banach space which is different from Hilbert spaces and which is isometric to subspaces of both L_p and L_q ?

It is a well-known fact first noticed by P.Levy that Hilbert spaces are isometric to subspaces of L_q for all $q > 0$. On the other hand, it was proved in [4] that, for $n \geq 3, q > 2, p > 0$, the function $\exp(-\|x\|_q^p)$ is not positive definite where $\|x\|_q = (|x_1|^q + \dots + |x_n|^q)^{1/q}$. (This result gave an answer to a question posed by I.J.Schoenberg [10] in 1938) In 1966, J.Bretagnolle, D.Dacunha-Castelle and J.L.Krivine [1] proved that, for $0 < p < 2$, a space $(E, \|\cdot\|)$ is isometric to a subspace of L_p if and only if the function $\exp(-\|x\|^p)$ is positive definite. Thus, in the language of isometries, the above mentioned result from [4] means that, for every $n \geq 3, q > 2, p \in (0, 2)$, the space l_n^q is not isometric to a subspace of L_p . (For $p \geq 1$, this fact was first proved in [2]) The initial purpose of this work was to find a non-Hilbertian subspace $(E, \|\cdot\|)$ of L_q with $q > 2$ of the dimension at least 3 such that the function $\exp(-\|x\|^p)$ is positive definite. The latter problem is equivalent to that at the beginning of the paper.

We prove, however, a more general fact: For every $n \geq 2$ and every compact subset Q of $(0, \infty) \setminus \{2k, k \in \mathbb{N}\}$, there exists an n -dimensional Banach space different from Hilbert spaces which is isometric to subspaces of L_q for all $q \in Q$ simultaneously.

In 1975, R.Schneider [9] proved that there exist non-trivial zonoids whose polars are zonoids or, in other words, there exist non-Hilbertian Banach spaces X such that X and X^* are isometric to subspaces of L_1 . It turns out that Schneider's construction of special subspaces of L_1 can be extended to all numbers $q > 0$ which are not even integers and in this way we obtain our main result.

2. Some properties of spherical harmonics.

We start with some properties of spherical harmonics (see [6] for details).

Let P_m denote the space of spherical harmonics of degree m on the unit sphere Ω_n in R^n . Remind that spherical harmonics of degree m are restrictions to the sphere of harmonic homogeneous polynomials of degree m . We consider spherical harmonics as functions from the space $L_2(\Omega_n)$. Any two spherical harmonics of different degrees are orthogonal in $L_2(\Omega_n)$ [6, p.2]

The dimension $N(n, m)$ of the space P_m can easily be calculated [6, p.4]:

$$(1) \quad N(n, m) = \frac{(2m + n - 2)\Gamma(n + m - 2)}{\Gamma(m + 1)\Gamma(n - 1)}$$

Let $\{Y_{mj} : j = 1, \dots, N(n, m)\}$ be an orthonormal basis of the space P_m . By the Addition Theorem [6, p.9], for every $x \in \Omega_n$,

$$(2) \quad \sum_{j=1}^{N(n, m)} Y_{mj}^2(x) = \frac{N(n, m)}{\omega_n}$$

where $\omega_n = 2\pi^{n/2}/\Gamma(n/2)$ is the surface of the sphere Ω_n .

Linear combinations of functions Y_{mj} are dense in the space $L_2(\Omega_n)$ [6, p.43]. Therefore, if F is a continuous function on Ω_n and $(F, Y_{mj}) = 0$ for every $m = 0, 1, 2, \dots$ and every $j = 1, \dots, N(n, m)$ then $F \equiv 0$ on Ω_n . Here (F, Y) stands for the scalar product in $L_2(\Omega_n)$.

Let Δ be the Laplace-Beltrami operator on the sphere Ω_n . Then for every $Y_m \in P_m$ we have [6, p.39]

$$(3) \quad \Delta Y_m + m(m + n - 2)Y_m \equiv 0$$

An immediate consequence of (3) (and a well-known fact) is that Δ is a symmetric operator and we can apply Green's formula: for every function H from the class C^{2r} , $r \in N$ of functions on Ω_n having continuous partial derivatives of order $2r$ and for every $Y_m \in P_m, m \geq 1$,

$$(4) \quad (-m(m + n - 2))^r (H, Y_m) = (H, \Delta^r Y_m) = (\Delta^r H, Y_m)$$

We also need the Funk-Henke formula [6, p.20]: for every $Y_m \in P_m$, every continuous function f on $[-1, 1]$ and every $x \in \Omega_n$,

$$(5) \quad \int_{\Omega_n} f(\langle x, \xi \rangle) Y_m(\xi) d\xi = \lambda_m Y_m(x)$$

where $\langle \cdot, \cdot \rangle$ stands for the scalar product in R^n and

$$(6) \quad \lambda_m = \frac{(-1)^m \pi^{(n-1)/2}}{2^{m-1} \Gamma(m + (n-1)/2)} \int_{-1}^1 f(t) \frac{d^m}{dt^m} (1 - t^2)^{m+(n-3)/2} dt$$

Let us calculate λ_m in the case where $f(t) = |t|^q, q > 0$.

LEMMA 1. If $q > 0, q \neq 2k, k \in N$ and $f(t) = |t|^q$ then

$$(7) \quad \lambda_m = \frac{\pi^{n/2-1} \Gamma(q+1) \sin(\pi(m-q)/2) \Gamma((m-q)/2)}{2^{q-1} \Gamma((m+n+q)/2)}$$

PROOF: Assume first that $q > m$ and calculate the integral from (6) by parts m times. Then use the formula $\int_{-1}^1 t^{2\alpha-1} (1-t^2)^{\beta-1} dt = \Gamma(\alpha) \Gamma(\beta) / \Gamma(\alpha + \beta)$ and formulas for Γ -function: $\Gamma(2x) = 2^{2x-1} \Gamma(x) \Gamma(x + 1/2) / \pi^{1/2}$ and $\Gamma(1-x) \Gamma(x) = \pi / \sin(\pi x)$. We get (7) for $q > m$. Note that both sides of (7) are analytic functions of q in the domain $\Re q > 0, q \neq 2k, k \in N$. Because of the uniqueness of analytic extension, (7) holds for every q from this domain. We are done. ■

3. Main result.

Let X be an n -dimensional subspace of $L_q = L_q([0, 1])$ with $q > 0$. Let $f_1, \dots, f_n$ be a basis in X and μ be the joint distribution of the functions $f_1, \dots, f_n$ with respect to Lebesgue measure (μ is a finite measure on R^n). Then, for every $x \in R^n$,

$$(11) \quad \|x\|^q = \left\| \sum_{k=1}^n x_k f_k \right\|^q = \int_0^1 \left| \sum_{k=1}^n x_k f_k(t) \right|^q dt = \int_{R^n} |\langle x, \xi \rangle|^q d\mu(\xi) = \int_{\Omega_n} |\langle x, \xi \rangle|^q d\nu(\xi)$$

where ν is the projection of μ to the sphere. (For every Borel subset A of Ω_n , $\nu(A) = \int_{\{tA, t \in R\}} \|x\|_2^q d\mu(x)$) The representation (11) of the norm is usually called the Levy representation. It is clear now that a norm in an n -dimensional Banach space admits the Levy representation with a probability measure on the sphere if and only if this space is isometric to a subspace of L_q . (Given the Levy representation we can choose functions $f_1, \dots, f_n$ on $[0, 1]$ with the joint distribution ν and define an isometry by $x \rightarrow \sum_{k=1}^n x_k f_k, x \in R^n$)

If we replace the measure ν by an arbitrary continuous (not necessarily non-negative) function on the sphere then a representation similar to the Levy representation is possible for a large class of Banach spaces (see [5] for the Levy representation with distributions instead of measures; such representation is possible for any Banach space and any q which is not an even integer) This is an idea going back to W.Blaschke that any smooth enough function on the sphere can be represented in the form (11) with a continuous function instead of a measure on the sphere. However, W.Blaschke and then R.Schneider [8] restricted themselves to the case $q = 1$ which is particularly important in the theory of convex bodies. The following theorem is an extension of R.Schneider's results from [8, p.77] and [9, p.367] to all positive numbers q which are not even integers

THEOREM 1. Let $q > 0, q \neq 2k, k \in N$ and let H be an even function of the class C^{2r} on Ω_n where $r \in N$ and $2r > n + q$. Then there exists a continuous function b_H on the sphere

Ω_n such that, for every $x \in \Omega_n$,

$$(8) \quad H(x) = \int_{\Omega_n} |\langle x, \xi \rangle|^q b_H(\xi) d\xi$$

Besides that, there exist constants $K(q)$ and $L(q)$ depending on n and q only such that, for every $x \in \Omega_n$,

$$(9) \quad |b_H(x)| \leq K(q) \|H\|_{L_2(\Omega_n)} + L(q) \|\Delta^{2r} H\|_{L_2(\Omega_n)}$$

PROOF: Define a function b_H on Ω_n by

$$(10) \quad b_H(x) = \sum_{m=0}^{\infty} \lambda_m^{-1} \sum_{j=1}^{N(n,m)} (N, Y_{mj}) Y_{mj}(x)$$

Since H is an even function and Y_{mj} are odd functions if m is odd the sum is, actually, taken over even integers m only.

Let us prove that the series in the right-hand side of (10) converges uniformly on Ω_n . By the Cauchy-Schwartz inequality, (2) and the fact that Y_{mj} form an orthonormal basis in P_m , we get

$$\begin{aligned} \left| \sum_{j=1}^{N(n,m)} (\Delta^r N, Y_{mj}) Y_{mj}(x) \right| &\leq \left(\sum_{j=1}^{N(n,m)} (\Delta^r N, Y_{mj})^2 \right)^{1/2} \left(\sum_{j=1}^{N(n,m)} Y_{mj}^2(x) \right)^{1/2} \\ &\leq \|\Delta^r H\|_{L_2(\Omega_n)} \left(\frac{N(n,m)}{\omega_n} \right)^{1/2} \end{aligned}$$

It follows from (4) and the latter inequality that

$$\begin{aligned} |b_H(x)| &\leq |\lambda_0^{-1} (H, Y_0) Y_0(x)| + \sum_{m=2; 2|m}^{\infty} \lambda_m^{-1} \left(\frac{-1}{m(m+n-2)} \right)^r \left| \sum_{j=1}^{N(n,m)} (\Delta^r H, Y_{mj}) Y_{mj}(x) \right| \leq \\ &|\lambda_0|^{-1} \omega_n^{-1/2} \|H\|_{L_2(\Omega_n)} + \sum_{m=2; 2|m}^{\infty} \lambda_m^{-1} m^{-2r} \left(\frac{N(n,m)}{\omega_n} \right)^{1/2} \|\Delta^r H\|_{L_2(\Omega_n)} \end{aligned}$$

Let us show that the series $\sum_{m=2; 2|m}^{\infty} \lambda_m^{-1} m^{-2r} (N(n,m)/\omega_n)^{1/2}$ converges. In fact, it follows from (1) that $N(n,m) = O(m^{n-2})$ and it is an easy consequence of (7) and the Stirling formula that $\lambda_m^{-1} = O(m^{(n+2q)/2})$. Since $2r > n + q = (n + 2q)/2 + (n - 1)/2 + 1$ we get $\lambda_m^{-1} m^{-2r} (N(n,m)/\omega_n)^{1/2} = o(m^{-1-\epsilon})$ for some $\epsilon > 0$, and the series is convergent. We denote the sum of this series by $L(q)$ and put $K(q) = |\lambda_0|^{-1} \omega_n^{-1/2}$, so we get (9).

We have proved that the series in (10) converges uniformly and defines a continuous function on Ω_n . It follows from (5) and the fact that all functions Y_{mj} are orthogonal that $(H, Y_{mj}) = (\int_{\Omega_n} |\langle x, \xi \rangle|^q b_H(\xi) d\xi, Y_{mj}(x))$ for every $m = 0, 1, 2, \dots$ and $j = 1, \dots, N(n, m)$. Hence, the function b_H satisfies (8). ■

Let X be an n -dimensional Banach space, $q > 0, q \neq 2k, k \in N$. Let $c(q) = \Gamma((n+q)/2)/(2\Gamma((q+1)/2)\pi^{(n-1)/2})$ be a constant such that $1 = c(q) \int_{\Omega_n} |\langle x, \xi \rangle|^q d\xi$ for every $x \in \Omega_n$. (The latter integral does not depend on the choice of $x \in \Omega_n$; it means that the norm of the space l_2^n admits the Levy representation with the uniform measure on the sphere and the space l_2^n is isometric to a subspace of L_q for every q)

Denote by $H(x), x \in \Omega_n$ the restriction of the function $\|x\|^q$ to the sphere Ω_n . Assume that the function H belongs to the class C^{2r} on Ω_n where $2r > n+q, r \in N$. Let b_H be the function corresponding to H by Theorem 1.

LEMMA 2. *If the number $K(q)\|H - 1\|_{L_2(\Omega_n)} + L(q)\|\Delta^r H\|_{L_2(\Omega_n)}$ is less than $c(q)$ then the space X is isometric to a subspace of L_q .*

PROOF: By (8) and definition of the number $c(q)$,

$$H(x) - 1 = \int_{\Omega_n} |\langle x, \xi \rangle|^q (b_H(\xi) - c(q)) d\xi$$

for every $x \in \Omega_n$. By (9), $|b_H(x) - c(q)| < c(q)$ for every $x \in \Omega_n$. It means that the function b_H is positive on the sphere. The equality (8) means that the space X admits the Levy representation with a non-negative measure and, by the reasoning at the beginning of Section 3, X is isometric to a subspace of L_q . ■

Now we are able to prove the main result of this paper. Let us only note that, for every function f of the class C^2 on the sphere Ω_n and for a small enough number λ , the function $N(x) = 1 + \lambda f(x), x \in \Omega_n$ is the restriction to the sphere of some norm in R^n . This is an easy consequence of the following one-dimensional fact: If $a, b \in R$, g is a convex function on $[a, b]$ with $g'' > \delta > 0$ on $[a, b]$ for some δ and $h \in C^2[a, b]$ then functions $g + \lambda h$ have positive second derivatives on $[a, b]$ for sufficiently small λ 's, and, hence, are convex on $[a, b]$.

THEOREM 2. *Let Q be a compact subset of $(0, \infty) \setminus \{2k, k \in N\}$. Then there exists a Banach space different from Hilbert spaces which is isometric to a subspace of L_q for every $q \in Q$.*

PROOF: Let f be any infinitely differentiable function on Ω_n and fix a number $r \in N$ so that $2r > n+q$ for every $q \in Q$. Choose a sufficiently small number λ such that the function $N(x) = 1 + \lambda f(x), x \in \Omega_n$ is the restriction to the sphere of some norm in R^n (see the

remark before Theorem 2) and such that, for every $q \in Q$, the function $H(x) = (N(x))^q$ satisfies the condition of Lemma 2. The possibility of such choice of λ follows from the facts that $K(q)$, $L(q)$ and $c(q)$ are continuous functions of q on the set Q and that $\|H - 1\|_{L_2(\Omega_n)}$ and $\|\Delta^{2r} H\|_{L_2(\Omega_n)}$ tend to zero uniformly with respect to $q \in Q$ as λ tends to zero. Now we can apply Lemma 2 to complete the proof. ■

Finally, let us consider the case where q is an even integer. It is easy to see that, for any fixed number $2k, k \in N, k > 1$, we can make the space X constructed in Theorem 2 isometric to a subspace of L_{2k} . In fact, let $N(x) = (1 + \lambda(x_1^{2k} + \dots + x_n^{2k}))^{1/4}$. For sufficiently small numbers λ , N is the restriction to the sphere of some norm in R^n and the corresponding space X is isometric to a subspace of L_q for every $q \in Q$. On the other hand, X is isometric to a subspace of L_{2k} because the norm admits the Levy representation with a measure on the sphere:

$$1 + \lambda(x_1^{2k} + \dots + x_n^{2k}) = \int_{\Omega_n} |\langle x, \xi \rangle|^{2k} (c(2k)d\xi + \lambda d\delta_1(\xi) + \dots + \lambda d\delta_n(\xi))$$

where δ_i is a unit mass at the point $\xi \in R_n$ with $\xi_i = 1, \xi_j = 0, j \neq i$.

Let us show that one can not make the space X isometric to subspaces of L_{2p} and L_{2q} if $p, q \in N$ and do not have common factors. In fact, if $(X, \|\cdot\|)$ is such a space then, for every $x \in R_n$,

$$\|x\|^{4pq} = \left(\int_{\Omega_n} |\langle x, \xi \rangle|^{2p} d\mu(\xi) \right)^{2q} = \left(\int_{\Omega_n} |\langle x, \xi \rangle|^{2q} d\nu(\xi) \right)^{2p}$$

for some measures μ, ν on Ω_n . The functions in the latter equality are polynomials and, since the polynomial ring has the unique factorization property, we conclude that $\|x\|^2$ is a homogeneous polynomial of the second order and X is a Hilbert space.

The situation is not clear if p and q have common factors. One can find some interesting results on Banach spaces with polynomial norms and on the structure of subspaces of $L_{2k}, k \in N$ in the paper [7].

Acknowledgements. I wish to thank Prof. Nigel Kalton for valuable remarks and helpful discussions during the work on this problem. I am grateful to Prof. Hermann König for bringing the paper [9] to my attention.

References.

1. Bretagnolle, J., Dacunha-Castelle, D. and Krivine, J.L.: *Lois stables et espaces L_p* , Ann. Inst. H.Poincare, Ser.B 2(1966), 231-259.
2. Dor, L.: *Potentials and isometric embeddings in L_1* , Israel J. Math. 24 (1976), 260-268.
3. Grzaslewicz, R.: *Plane sections of the unit ball of L_p* , Acta Math. Hung. 52(1988), 219-225.
4. Koldobsky, A.: *Schoenberg's problem on positive definite functions*, Algebra and Analysis 3 (1991), No. 3, 78-85 (Russian); English translation in St.Petersburg Math. J. 3 (1992), 563-570
5. Koldobsky, A.: *Generalized Levy representation of norms and isometric embeddings into L_p -spaces*, Ann. Inst. H.Poincare (Prob. and Stat.) 28 (1992), 335-353
6. Muller, C.: *Spherical Harmonics*, Lect. Notes in Math. 17, Springer-Verlag, Berlin, 1966.
7. Reznick, B.: *Banach spaces with polynomial norms*, Pacific J. Math. 82 (1979), 223-235.
8. Schneider, R.: *Zu einem problem von Shephard uber die projektionen konvexer korper*, Math. Zeitschrift 101 (1967), 71-82.
9. Schneider, R.: *Zonoids whose polars are zonoids*, Proc. Amer. Math. Soc. 50 (1975), 365-368.
10. Schoenberg, I.J.: *Metric spaces and positive definite functions*, Trans. Amer. Math. Soc. 44 (1938), 522-536.